arXiv:1304.5836v1 [q-bio.MN] 22 Apr 2013

Oscillation in microRNA Feedback Loop

Bin Ao, Sheng Zhang, Caiyong Ye, Lei Chang, Guangming Zhou, and Lei Yang[1]
Institute of Modern Physics, Chinese Academy of Science, Lanzhou 730000, China

(Dated: April 23, 2013)

The dynamic behaviors of microRNA and mRNA under external stress are studied with biological experiments and mathematics models. In this study, we developed a mathematic model to describe the biological phenomenon and for the first time reported that, as responses to external stress, the expression levels of microRNA and mRNA sustained oscillation. And the period of the oscillation is much shorter than several reported transcriptional regulation negative feedback loop.

PACS numbers:

Since the first microRNA (miRNA) lin-4 was discovered two decades ago, microRNA has been a rising star in the field of biology. It has been proved to be related with lots of essential processes in cells, such as DNA damage response, cell-cycle checkpoints, programmed cell death and metabolism [2]. As known to all, microRNA executes its functions as a regulator by binding to complementary sequences on target messenger RNA transcripts (mRNA) and forming RNA-induce silencing complex (RISC), after transcription but before translation. Length of a mature miRNA is just about 22 nt and a gene-coding mRNA is much bigger and needs to translate into a protein. Therefore, comparing with the regulation by a protein, the regulatory function of a miRNA is a more economical means [2].

In cell signaling pathway, negative feedback loop is a very normal motif, consisted of not only proteins but also other non-coding RNAs including miRNAs. The advantage of this topological mode lies in its stability [3]. There are many researches on protein-protein negative feedback loop, e.g., mdm2 and p53 [4–6], but few on miRNA regulations [7–10]. Based on the special biological functions of miRNAs, we built a stress-induced miRNA-protein dynamical model which contains a negative feedback loop. Then we designed a series of experiments to observe dynamic behaviors of this regulation on molecular level. We made a first report on miRNA oscillation in cells exposed to radiation and these results generally support our model. Furthermore, this model could be used to construct dynamical networks of proteins and other non-coding RNAs.

A scheme of miRNA feedback loop is shown in Fig. 1(a). The mRNA transcription of a certain gene increases under stress, and its corresponding protein, as a transcriptional factor, actives the expression of the miRNA which represses the transcription of the mRNA in return. A three-variable ODEs is considered to describe the system,

$$\begin{aligned}\frac{d[mRNA]}{dt} &= Sig - k_{mRNA-protein}[mRNA]\\ &-d_{mRNA}[mRNA] - k_{miRNA-mRNA}\frac{[miRNA][mRNA]}{j_{mim}+[mRNA]},\\[1ex] \frac{d[protein]}{dt} &= k_{mRNA-protein}[mRNA] - d_{protein}[protein],\\ \frac{d[miRNA]}{dt} &= k_{protein-miRNA}[protein] - d_{miRNA}[miRNA],\end{aligned} \tag{1}$$



FIG. 1: (a) A scheme of miRNA feedback loop. A signal of radiation induces transcription of mRNA, and its corresponding protein, as a transcriptional factor, active the transcription miRNA. Then the miRNA inhibits the expression of protein by inducing the degradation of mRNA. (b) A scheme of p53-mdm2 negative feedback loop.

where $[mRNA]$, $[protein]$, $[miRNA]$ are the expression levels of the mRNA, protein, and miRNA, respectively. And Sig is the signal that stimulates the responding effects of cells and activates the transcription of mRNA. For simplicity, Sig is just a constant here. As will be shown below, a non-zero Signal is a necessary condition of the excitation of a sustained oscillation. d_{mRNA}, $d_{protein}$, and d_{miRNA} are degradation rates of the three molecules. $k_{mRNA-protein}$ and $k_{protein-miRNA}$ are the rates of protein translation and miRNA transcription. Only a linear model is considered. Finally, for the regulation of miRNA and mRNA, as RISC is a stable machine [2], we tend to consider the condition that miRNA could be recycled, and then the Michael-Mathen function is considered. By renormalizing time and parameters according to

$$\begin{gathered}\tau = d_{miRNA}t, S = Sig/d_{miRNA},\\ d_m = (d_{mRNA} + k_{mRNA-protein})/d_{miRNA},\\ k_{mim} = k_{miRNA-mRNA}/d_{miRNA},\\ d_p = d_{protein}/d_{miRNA},\\ k_{mp} = k_{mRNA-protein}/d_{miRNA},\\ d_mi = d_{miRNA}/d_{miRNA} = 1,\\ k_{pmi} = k_{protein-miRNA}/d_{miRNA},\end{gathered} \tag{2}$$

Eq. (1) can be rewritten in a simplified form,

$$\begin{aligned} \dot{m} &= S - d_m m - k_{mim}\frac{m_i m}{j_{mim}+m}, \\ \dot{p} &= -d_p p + k_{mp} m, \\ \dot{m_i} &= -m_i + k_{pmi} p. \end{aligned} \tag{3}$$

The diurnal oscillation of miRNA and mRNA are reported in recent works [11]. In this paper, we report that this phenomenon can also be observed in stress-response process. And the dynamical behaviors of Eq. (3) with proper parameters shows sustained oscillation, which can be used to describe the biological phenomenon. Fig. 2 shows an example. Moreover, it can also be observed that the periods of the three oscillation curves are almost the same, but their phases are different. The oscillation of mRNA is firstly excited by a non-zero signal, and followed by the oscillation of the protein. And then the expression level of miRNA also rises and begins to repress the production of mRNA. Thus the oscillations of mRNA and miRNA are almost anti-phase.

Since there have been many studies on transcription, translation and the degradation of mRNA and protein, and miRNA regulation negative feedback loop can be analogous to the well studied p53-mdm2 feedback loop shown in Fig. 1(b), the parameters about these processes are easy to be chosen [12]. But the transcription and degradation rates of miRNA and the parameters of miRNA-mRNA interaction are still unclear. Thus, these parameters will be mainly focused on in this paper. Since the degradation rate of miRNA (d_{mi}) has been reduced, only are three parameters (k_{mim}, k_{pmi}, j_{mim}) left.

Because all parameters are positive, system (3) has only one positive fixed point. And as will be shown below, $0 \approx j_{mim} << k_{mim}, k_{pmi}$. Thus the fixed point can be written by

$$\begin{aligned} m^* &= \frac{9 * k_{pmi}}{8 + 10 * k_{mim} * k_{pmi}}, \\ p^* &= \frac{9}{8 + 10 * k_{mim} * k_{pmi}}, \\ m_i^* &= \frac{9}{8 + 10 * k_{mim} * k_{pmi}}. \end{aligned} \tag{4}$$

To assess whether Eq. (3) can display oscillations, we have searched for parameter values where the fixed point specified by (4) loses stability to a periodic solution via a Hopf bifurcation, that is, a pair of conjugate eigenvalues of the linearized problem cross the imaginary axis. Because Eq. (3) has three variables, the characteristic polynomial of Jacobi matrix in the fixed point (4) is a cubic monic polynomial, which is denoted by

$$\begin{aligned} p(\lambda) &= \lambda^3 + b\lambda^2 + c\lambda + d \\ b &= \frac{65m^{*2} + 25k_{mim}m_i^* j_{mim}}{25m^{*2}} \\ c &= \frac{56m^{*2} + 45k_{mim}m_i^* j_{mim}}{25m^{*2}} \\ d &= \frac{4(4+5k_{mim}k_{pmi})m^{*2} + 20(m_i^* - k_{pmi}m^*)k_{mim}j_{mim}}{25m^{*2}}. \end{aligned} \tag{5}$$



FIG. 2: The time course of the expression of mRNA (green line), protein (red line) and miRNA (blue line) described by Eq. (3) with $S = 0.9$, $dm = 0.8$, $kmim = 6$, $jmim = 0.001$, $dp = 0.8$, $kmp = 0.8$ $dmi = 3$, $kmpi = 10$, and $mi = 0.135$, $m = 0.047$, $p = 0.028$ at $t = 0$.

The roots of Eq. (5) are the eigenvalues of the linearized form of Eq. (3). They describe the dynamics behavior of Eq. (3) around the fixed point. If the determinant of Eq. (5)

$$\Delta = (\frac{bc}{6} - \frac{b^3}{27} - \frac{d}{2})^2 + (\frac{c}{3} - \frac{b^2}{9})^3 \tag{6}$$

is positive, the linearized Eq. (3) will has one real eigenvalue (denoted by λ_1) and two conjugate imaginary eigenvalues (λ_2 and λ_3). That is, the oscillation of system Eq. (3) will be excited around the fixed point Eq. (4). But, a sustained oscillation requires that the real parts of λ_2 and λ_3 cross 0, i.e., the Hopf bifurcation. Thus according to the general formula of the roots of Eq. (5), the critical point of Hopf bifurcation can be written as

$$-\frac{b}{3} - \frac{1}{2}(\sqrt[3]{T + \sqrt{T^2 + K^3}} + \sqrt[3]{T - \sqrt{T^2 + K^3}}) = 0. \tag{7}$$

where $T = bc/6 - b^3/27 - d/2$ and $K = c/3 - b^2/9$. Then, to cubing both sides of Eq. (7), it will be

$$-\frac{8b^3}{27} = 2T - 2bK. \tag{8}$$

Finally, we have

$$R = d - bc = 0. \tag{9}$$

That is, the condition of the existence of a sustained oscillation in Eq. (3) is

$$R > 0. \tag{10}$$

Thus, to substitute the parameters of Eq. (3) into Eq. (6) and Eq. (10) and to solve the equation $R = 0$,one can obtain the phase graph in parameter space (k_{mim}, j_{mim}, k_{pmi}).

It is easy to be tested that, no matter in k_{mim}-k_{pmi} plane or in k_{mim}-j_{mim} plane, the determinant Eq. (6) has



FIG. 3: The attenuate oscillation area (labeled by A) and the sustained oscillation area (labeled by S) in (a) k_mim-k_pmi and (b) j_mim-k_mim parameter space, which are divided by the critical line satisfied $R = 0$.



FIG. 4: (a) The m_i curves of Eq. (3) with 1000 different parameter values. (b) m_i, m, p curves of Eq. (3) with $S = 0.9$, $dm = 0.8$,$kmim = 6$, $jmim = 0.001$, $dp = 0.8$, $kmp = 0.8701$ $dmi = 3$, $kmpi = 7.3298$, and $mi = 0.135$, $m = 0.047$, $p = 0.028$ at $t = 0$. (c) The curves of $< m >$, $< p >$ and $< m_i >$

positive values in a broad parameter area. That means an oscillation solution of the system (3) is usual and easy to be obtained. But, a sustained oscillation is unusual. Firstly, the oscillation parameter area($R > 0$) in k_{mim}-k_{pmi} plane is shown in Fig. 3(a). This picture can be explained in the sense that an over faster transcription rate of miRNA (k_{pmi} is big) and an over powerful repression to mRNA (k_{mim} is big simultaneously) will dramatically reduce the production of mRNA and protein, and then oscillation attenuates. Furthermore, in k_{mim}-j_{mim} plane (Fig. 3(b)), $R > 0$ can only be achieved when j_{mim} is very small. That is, Eq. (3) is a weak nonlinear dynamics system. Since the Michael-Mathen coefficient j_{mim} is the ratio of the rate of a reaction and that of its inverse reaction, a very small j_{mim} means the reaction is basically irreversible. And this feature can be observed in miRNA-mRNA interaction procession.

Empirically speaking, the oscillation of miRNA and protein in a single cell will not reduce under a constant drive, but the peak could be broadened because of the non-synchronization among the cells. The model Eq. (3) only describes mono cell behavior. But in real biological systems, the rates of transcription, translation and other regulations vary in different cells. This will lead to different parameters when the model Eq. (3) is used to describe different cells. Thus, when the problem refers to the behaviors of millions of cells, the parameter robustness of the system Eq. (3) must be discussed. Fig. 4(a) shows the curves of the system Eq. (3) with different parameters. It can be found that the system Eq. (3) is parameter robust. Even though the environments can affect some specific biological processes, the existence of sustained oscillation will not be dramatically changed by small perturbation of environments. On the other hand, the amplitudes and periods of oscillations depend on parameters values. Therefore, the collective behavior of cells will show amplitude attenuation. To illustrate this picture, Eq. (3) is simulated by using N=10000 different parameter values (the expectation of these parameter values is equal to that we used before and their standard deviation is less than 30% of the values that we used before). Then the averages over the 10000 dynamical processes are calculated: $< m(t) >= \sum m^k(t)/10000$, $< p(t) >= \sum p^k(t)/10000$, $< m_i(t) >= \sum m_i^k(t)/10000$, where k is the index of different parameter values. As shown in Fig. 4(b) and Fig. 4(c), the oscillation of $< m >$, $< p >$ and $< m_i >$ attenuate.



FIG. 5: The experimental data of Dicer mRNA (green line), Dicer protein (red line) and miRNA-3928 (blue line) and the analysis results of model Eq. (3).

To testify the theoretical result, an experiment is designed. We chose human cervical carcinoma HeLa cell line for the biological experiment. Dicer protein is required to process miRNA precursor to mature miRNAs, and meanwhile is a target of miRNA-3928 . So Dicer and miRNA-3928 form a negative feedback loop as shown in Fig. 1(a). HeLa cells were cultured in RPMI-1640 media and exposed to 2 Gy X-rays at room temperature. The qRT-PCR (quantitative real-time reverse transcription-PCR) is used to measure time-sequencing expression levels of miRNA-3928 and Dicer mRNA. After irradiation, DNA repair were activated in cells. Unless the damages



FIG. 6: Western blotting picture of Dicer protein till 2.25 hours after radiation, in which the oscillation of its concentration can be found. And the oscillation period is about 1 hour.

are repaired, cells would not reenter cell cycle. Under the conditions of dose used in this experiment, the duration of repair is about 6-8 hours. The drive would disappear at that time [14]. The results are shown by the circle points in Fig. 5. And Western blotting is used to measure the expression level of Dicer protein. The concentration of Dicer is shown by circle points in Fig. 5, and the oscillation of Dicer protein can also be found in one of Western blotting pictures shown in Fig. 6. To get the numerical data, Gray value analysis is used to process pictures. And it can be found that the concentrations of miRNA-3928, Dicer and its mRNA show an obvious oscillation behavior after radiation. The oscillation period is about 1 hour, and the amplitudes of the oscillations attenuate after several oscillation periods. The phases of these oscillations are different, opposite change trend between the concentration of Dicer and that of miRNA-3928 can always be observed.

On the other hand, the transcriptional regulation is also popular in biological processes. The oscillations of proteins p53 and mdm2 with a phase differences can also be observed and various mathematical models were constructed to explain the experiments. But a significant difference between transcriptional regulation and miRNA regulation is the oscillation periods. The period of p53 and mdm2 oscillation is about 6 hours, while that of miRNA-3928 and Dicer oscillation is only 1 hour. This shows that the response time of miRNA regulation is far more fasten than that of transcriptional factor regulation. Obviously, a faster response is very important to radiation response, particularly to the repairs of radiation damages.

In conclusion, we build a model to describe the miRNA-protein negative feedback loop, which is tested by the biology experiment of Dicer and miRNA-3928 regulation loop. Through the numerical simulation and experiments, the oscillations of the expression of mRNA, protein and miRNA and their phase difference are observed. We find that the response time of miRNA regulation is far faster than that of the transcriptional factor regulation. The results of simulations are consistent with those of experiments well. These results provide estimates of the effective biochemical parameter in the processes involving miRNA and could be used to construct dynamical networks of proteins and other noncoding RNAs.

---

[1] Corresponding author: lyang@impcas.ac.cn

[2] J. Krol, I. Loedige, W. Filipowicz, Nature Reviews Genetics, **11**, 597-610 (2010).

[3] Uri Alon, *An Introduction to Systems Biology: Design Principles of Biological Circuits*, (CRC Press, 2006)

[4] Ruth Lev Bar-Or, Ruth Maya, Lee A. Segel, Uri Alon, Arnold J. Levine, and Moshe Oren, PNAS, **97(21)**, 11250-11255 (2000).

[5] Xiao-Peng Zhang, Feng Liu, Zhang Cheng, Wei Wang, PNAS, **106(30)**, 12245-12250 (2009)

[6] Naama Geva-Zatorsky, Erez Dekel, Eric Batchelor, Galit Lahav, Uri Alon, PNAS, **107(30)**, 13550-13555 (2010)

[7] J. A. Pulikkan et al., Blood, **115(9)**, 1768-1778 (2010).

[8] S. Tokumaru et al, Carcinogenesis, **29(11)**, 2073-2077 (2008).

[9] Peipei Zhou, Shuiming Cai1, Zengrong Liu, and Ruiqi Wang, Physical Review E, **85**, 041916 (2012).

[10] Yichen Li,Yumin Li mail, Hui Zhang, Yong Chen, PLoS ONE, **6(10)**: e26302.

[11] C. Sire, A. B. Moreno, M. Garcia-Chapa, J. J. Lpez-Moya, B. S. Segundo, FEBS letters, **583(6)**, 1039-1044(2009).

[12]

[13] Aziz SANCAR, Laura A. LINDSEY-BOLTZ, Keziban NSAL-KACMAZ, Stuart LINN, Annual review of biochemistry, **73**, 39-85(2004).

[14] Naama Geva-Zatorsky, Nitzan Rosenfeld, Shalev Itzkovitz, Ron Milo, Alex Sigal, Erez Dekel, Talia Yarnitzky, Yuvalal Liron,Paz Polak, Galit Lahav and Uri Alon, Molecular Systems Biology, 2006.0033.